

LINEARITY OF HOMOGENEOUS ORDER ONE SOLUTIONS TO ELLIPTIC EQUATIONS IN DIMENSION THREE

QING HAN, NIKOLAI NADIRASHVILI, AND YU YUAN

1. Introduction

In this note we prove that any homogeneous order one solution to non-divergence elliptic equations in $\mathbb{R}^3$ must be linear. Consider the general equations in $\mathbb{R}^n$

$$(1.1) \quad \sum_{i,j=1}^n a_{ij}(x) D_{ij} u = 0,$$

where the coefficients satisfy

$$\lambda I \leq (a_{ij}(x)) \leq \lambda^{-1} I$$

for some positive constant $\lambda > 0$, and the dimension $n \geq 3$. Safonov constructed homogeneous order α solutions, with $\alpha \in (0, 1)$, to (1.1) in [10], where he showed the unimprovability of the estimates of Hölder exponent for solutions to (1.1) by Krylov and himself. The homogeneity α with $\alpha < 1$ plays an essential role in the construction. Later on, Safonov asked in [11, p.49] whether one can construct nontrivial homogeneous order one solutions to (1.1). Our result indicates that it is impossible to do so in $\mathbb{R}^3$.

Theorem 1.1. *Any homogeneous order one strong solution u to (1.1) in $\mathbb{R}^3$ with $u \in W_{loc}^{2,2}(\mathbb{R}^3)$ must be a linear function.*

On the other hand, one does have nontrivial homogeneous order one solutions to (1.1) in $\mathbb{R}^4 \setminus \{0\}$. In fact let $(x, f(x)) \in \mathbb{R}^n \times \mathbb{R}^k$ be a nonparametric minimal surface. Then each component of f satisfies (1.1), with $(a_{ij}(x))$ being the inverse of the induced metric of the minimal surface in $\mathbb{R}^{n+k}$ (cf. [7, p.3]). Through Hopf fibration, Lawson and Osserman constructed a minimal cone $(x, f(x)) \in \mathbb{R}^4 \times \mathbb{R}^3$, where

$$f(x) = \frac{\sqrt{5}}{2} \frac{1}{|x|} (x_1^2 + x_2^2 - x_3^2 - x_4^2, 2x_1x_3 + 2x_2x_4, 2x_2x_3 - 2x_1x_4)$$

(see [7, Theorem 7.1]). Now each component of f is a desired nontrivial solution to (1.1) in $\mathbb{R}^4 \setminus \{0\}$. Actually by noticing that the graph of

Date: October 29, 2001. Submitted on November 12, 2001.

All authors are partially supported by NSF grants, and the first author also by Sloan Research Fellowship.

$u = (x_1^2 + x_2^2 - x_3^2 - x_4^2) / |x|$ is a saddle surface, one can easily construct coefficients $a_{ij}(x)$ in $\mathbb{R}^4 \setminus \{0\}$ so that u satisfies (1.1) in $\mathbb{R}^4 \setminus \{0\}$.

Theorem 1.1 gives a simple ‘‘PDE’’ proof of a well-known result obtained by many authors in 1970’s, which states that any nonparametric minimal cone of dimension three must be flat. Let $(x, f(x)) \in \mathbb{R}^3 \times \mathbb{R}^k$ be the minimal cone with $f(tx) = tf(x)$, $f \in C^\infty(\mathbb{R}^3 \setminus \{0\})$, then each component of f satisfies (1.1). By Theorem 1.1, f must be linear, or the minimal cone is flat.

From Theorem 1.1 one also sees that any smooth homogeneous order two solution in $\mathbb{R}^3 \setminus \{0\}$ to the fully nonlinear elliptic equation $F(D^2u) = 0$ must be a quadratic polynomial. To get this conclusion, one simply apply Theorem 1.1 to the gradient ∇u and (1.1) with $a_{ij}(x) = \frac{\partial F}{\partial m_{ij}}(D^2u)$. In contrast, the second author constructed a non-quadratic homogeneous order two solution to some equation $F(D^2u) = 0$ in $\mathbb{R}^{12}$, which provides a counterexample to the regularity for fully nonlinear elliptic equations (see [8]).

The heuristic idea of the proof of Theorem 1.1 is the following simple geometric observation. The closed saddle surface $\nabla u(\mathbb{S}^2)$ must be a point. Therefore, u is linear. More precisely, one consider the surface Σ parametrized by the gradient $\nabla u : \mathbb{S}^2 \rightarrow \mathbb{R}^3$. Because of (1.1), Σ is a saddle surface at $\nabla u(x)$ with $D^2u(x) \neq 0$. However the supporting plane with normal x touches Σ at $\nabla u(x)$. Thus $D^2u(x) \equiv 0$ and it follows that u is linear, see Section 2.

The ideas of gradient map and supporting plane are already in an early paper by Alexandrov [1]. In fact, under the assumption that the homogeneous order one function u is analytic in $\mathbb{R}^3 \setminus \{0\}$ and the Hessian D^2u is either non-definite or 0 at each point, Alexandrov showed that u must be a linear function. Roughly speaking, if u is analytic in $\mathbb{R}^3 \setminus \{0\}$, the set $\mathcal{S}(u) = \{x \in \mathbb{S}^2 | D^2u(x) = 0\}$ is either isolated or the whole $\mathbb{S}^2$. Alexandrov proved that the supporting plane to Σ is unique at $\nabla u(x)$ with x being an isolated point of $\mathcal{S}(u)$. Since the surface Σ has supporting planes with normal along all the directions in $\mathbb{R}^3$, Alexandrov excluded case of $\mathcal{S}(u)$ being isolated and proved the result. It is interesting to note that the concept of gradient maps and supporting planes was further employed in the later development of the Alexandrov-Bakel’man-Pucci maximum principle.

When the coefficients a_{ij} are C^α and the solution u is $C^{2,\alpha}$ away from the origin, we can show that the set $\mathcal{S}(u)$ is either isolated or the whole $\mathbb{S}^2$. Coupled with Alexandrov’s argument, one sees that Theorem 1.1 holds in the $C^{2,\alpha}$ setting. After the work in $C^{2,\alpha}$ case was done, we found that Pogorelov [9] generalized the above Alexandrov’s result to C^2 functions. The argument in [9] is more involved. Since our approach in $C^{2,\alpha}$ case is interesting in its own right and short, we also include it here in Section 3.

The authors wish to thank L. Nirenberg for informing and mailing them the inspiring reference [1].

2. Proof of the Main Theorem

Let $h(x_1, x_2) = u(x_1, x_2, 1)$. Then the homogeneous order one function $u(x_1, x_2, x_3) = x_3 h\left(\frac{x_1}{x_3}, \frac{x_2}{x_3}\right)$, and the gradient ∇u and the Hessian $D^2 u$ have the following representation

$$(2.1) \quad \nabla u(x_1, x_2, 1) = (h_1, h_2, h - x_1 h_1 - x_2 h_2),$$

$$(2.2)$$

$$D^2 u(x_1, x_2, 1) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -x_1 & -x_2 & 1 \end{bmatrix} \begin{bmatrix} h_{11} & h_{12} & 0 \\ h_{21} & h_{22} & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & -x_1 \\ 0 & 1 & -x_2 \\ 0 & 0 & 1 \end{bmatrix}.$$

From our assumption that the homogeneous order one solution u is in $W_{loc}^{2,2}(\mathbb{R}^3)$, it follows that $h \in W_{loc}^{2,2}(\mathbb{R}^2)$ is a strong solution to

$$(2.3) \quad \sum_{i,j=1}^2 A_{ij}(x_1, x_2) D_{ij} h = 0,$$

where the coefficients A_{ij} are in terms of $a_{ij}(x_1, x_2, 1)$, x_1 , x_2 and satisfy the ellipticity condition with some $\lambda(x_1, x_2)$.

Lemma 2.1. *For any $\nu \in \mathbb{S}^2$, the supporting plane with normal ν must touch the surface Σ parametrized by $\nabla u : \mathbb{S}^2 \rightarrow \mathbb{R}^3$ at $\nabla u(\nu)$ or $\nabla u(-\nu)$.*

Proof. First we notice that $\nabla u \in C^\alpha(\mathbb{S}^2)$. In fact each component of (h_1, h_2) , say h_1 is a $W_{loc}^{1,2}(\mathbb{R}^2)$ weak solution to a divergence equation

$$\sum_{i,j=1}^2 D_i (B_{ij} D_j h) = 0,$$

where $B_{11} = A_{11}/A_{22}$, $B_{12} = 2A_{12}/A_{22}$, $B_{22} = 1$. It follows that $h \in C^{1,\alpha}(S^2)$ (cf. [5, p.284-285 or Theorem 12.4]). By (2.1) we see $\nabla u \in C^\alpha(\mathbb{S}^2)$ with α depending only on the original ellipticity λ . Therefore, for any $\nu \in \mathbb{S}^2$, the supporting plane P_ν with normal ν must touch the surface Σ somewhere (Σ is on the opposite side of ν).

Without loss of generality, we assume $\nu = (1, 0, 0)$.

Claim: $P_{(1,0,0)}$ touches Σ at $\nabla u(x_1, x_2, 0)$ with $(x_1, x_2, 0) \in \mathbb{S}^2$.

Suppose $P_{(1,0,0)}$ touches Σ at $\nabla u(x_1, x_2, x_3)$ with $x_3 \neq 0$, say $x_3 > 0$. Then $u_1(x_1, x_2, 1) = u_1\left(\frac{x_1}{x_3}, \frac{x_2}{x_3}, 1\right) = h_1\left(\frac{x_1}{x_3}, \frac{x_2}{x_3}\right)$ would achieve its maximum at $\left(\frac{x_1}{x_3}, \frac{x_2}{x_3}\right)$. Because of (2.3), h_1 satisfies the strong maximum principle (cf. [5, Theorem 8.19]). We then have a contradiction unless $h_1 \equiv \text{const.}$. In the latter case, $u_1(x_1, x_2, x_3) \equiv \text{const.}$ for $x_3 > 0$, our claim holds.

Similarly, by applying the above argument to $u(x_1, 1, x_3)$, we see that $P_{(1,0,0)}$ also touches Σ at $\nabla u(x_1, 0, x_3)$ with $(x_1, 0, x_3) \in \mathbb{S}^2$. Therefore, $P_{(1,0,0)}$ must touch Σ at $\nabla u(1, 0, 0)$ or $\nabla u(-1, 0, 0)$. $\square$

We now present the proof of Theorem 1.1.

Proof. By our assumption on u , $u \in W^{2,2}(\mathbb{S}^2)$. If $D^2u = 0$ almost everywhere on $\mathbb{S}^2$, then u is already linear. Otherwise, we pick a Lebesgue point $x^* \in \mathbb{S}^2$ for D^2u with $D^2u(x^*) \neq 0$, say $x^* = (1, 0, 0)$. We may also assume x^* is a Lebesgue point for ∇u and $(a_{ij}(x))$. By Lemma 2.1, the supporting planes $P_{(1,0,0)}$ and $P_{(-1,0,0)}$ touches Σ at $\nabla u(1, 0, 0)$ or $\nabla u(-1, 0, 0)$. If both planes touch Σ at the same point, then we see that $u_3 \equiv c$. Consequently the homogeneous order one function $v(x_1, x_2) = u - cx_3$ satisfies (1.1) and it follows that v is linear, or u is also linear.

So we are left with the case that $P_{(1,0,0)}$ and $P_{(-1,0,0)}$ touch Σ at different point. We may assume $P_{(1,0,0)}$ touches Σ at $\nabla u(1, 0, 0)$. It means that

$$(2.4) \quad u_3(x_1, x_2, 1) = h - x_1 h_1 - x_2 h_2 \leq h(0) \quad \text{near } (0, 0).$$

By (2.1) and (2.2), $(0, 0)$ is a Lebesgue point for D^2h and ∇h . It follows that (cf. [4, Appendix C])

$$h(x_1, x_2) = h(0) + h_1(0)x_1 + h_2(0)x_2 + \frac{1}{2} \sum_{i,j=1}^2 h_{ij}(0)x_i x_j + o(|x|^2).$$

Because of (2.3) and $D^2h(0, 0) \neq 0$, we may assume $h_{11}(0) = a$, $h_{12}(0) = 0$, $h_{22}(0) = -b$ with $a, b > 0$. A simple computation yields $u(0, x_2, 1) = h(0) + \frac{1}{2}bx_2^2 + o(x_2^2)$ which contradicts (2.4). This completes the proof. $\square$

3. Another Proof in the $C^{2,\alpha}$ Case

In this section, we present yet another proof of Theorem 1.1 in the $C^{2,\alpha}$ case. Suppose $u : \mathbb{R}^3 \setminus \{0\} \rightarrow \mathbb{R}$ is a $C^{2,\alpha}$ homogeneous order one function $u = rg(\theta_1, \theta_2)$, where (r, θ_1, θ_2) is the spherical coordinates with $x_1 = r \cos \theta_2 \cos \theta_1$, $x_2 = r \cos \theta_2 \sin \theta_1$, $x_3 = r \sin \theta_2$. Then the Hessian $D^2u = \left(\frac{\partial^2 u}{\partial x_i \partial x_j} \right)$ has the following representation

$$(3.1) \quad D^2u(x) = \frac{1}{r} \mathbf{R}(\theta_1, \theta_2) H \mathbf{R}'(\theta_1, \theta_2)$$

where

$$H = \begin{bmatrix} 0 & 0 & 0 \\ 0 & \frac{1}{\cos^2 \theta_2} \frac{\partial^2 g}{\partial \theta_1^2} - \tan \theta_2 \frac{\partial g}{\partial \theta_2} + g & \frac{\partial^2 g}{\partial \theta_1 \partial \theta_2} + \frac{\sin \theta_2}{\cos^2 \theta_2} \frac{\partial g}{\partial \theta_1} \\ 0 & * & \frac{\partial^2 g}{\partial \theta_2^2} + g \end{bmatrix}$$

and $\mathbf{R}(\theta_1, \theta_2)$ is the rotation from $(\partial x_1, \partial x_2, \partial x_3)$ to $(\partial r, \frac{1}{r} \partial \theta_1, \frac{1}{r} \partial \theta_2)$. We see that $D^2u = 0$ if and only if $H = 0$. We define the singular set $S(u)$ as

$$S(u) = \{x \in \mathbb{S}^2 \mid D^2u(x) = 0\}.$$

Lemma 3.1. *Let u be as in Theorem 1.1. Then one of the following holds:*

- (1) $S(u)$ is empty;
- (2) $S(u)$ consists of finitely many points;
- (3) $S(u) = \mathbb{S}^2$.

Proof. Suppose $S(u)$ is not empty. For any $p \in S(u)$, we prove that either p is isolated or $S(u)$ contains a neighborhood of p on $\mathbb{S}^2$. Without loss of generality, we assume $p = (1, 0, 0)$, $a_{ij}(p) = \delta_{ij}$. Note that $S(u) = S(u + ax_1 + bx_2 + cx_3)$, we may also assume $\nabla u(p) = 0$, then $u(p) = p \cdot \nabla u(p) = 0$. Correspondingly, we have

$$\left(\frac{\partial^2 g}{\partial \theta_i \partial \theta_j} \right) (0, 0) = 0, \quad \left(\frac{\partial g}{\partial \theta_1}, \frac{\partial g}{\partial \theta_2} \right) (0, 0) = 0, \quad \text{and} \quad g(0, 0) = 0.$$

Also g satisfies

$$\sum_{i,j=1}^2 A_{ij}(\theta_1, \theta_2) \frac{\partial^2 g}{\partial \theta_i \partial \theta_j} + \sum_{i=1}^2 B_i(\theta_1, \theta_2) \frac{\partial g}{\partial \theta_i} + C(\theta_1, \theta_2) g = 0,$$

where the C^α coefficients (A_{ij}) satisfy the ellipticity condition with the same constant λ , $A_{ij}(0, 0) = \delta_{ij}$, B_i and C are C^α functions near $(0, 0)$. Suppose g vanishes up to infinity order at $(0, 0)$, then by the Carleman unique continuation (cf. [3, p.124]), $g \equiv 0$ in a neighborhood of $(0, 0)$. It follows that $D^2 u = 0$ on $\mathbb{S}^2$, since u is C^2 . Suppose g vanishes up to order $k-1$ at $(0, 0)$. By our assumption $k \geq 3$. We apply the result of [2, Theorem 1] to obtain

$$g = P(\theta_1, \theta_2) + R(\theta_1, \theta_2),$$

where the homogeneous order k polynomial P satisfies $\sum_{i,j=1}^2 A_{ij}(0, 0) \frac{\partial^2 P}{\partial \theta_i \partial \theta_j} = 0$, or $\Delta P = 0$, and the remainder satisfies

$$R \sim O(|\theta|^{k+\alpha}), \quad \nabla R \sim O(|\theta|^{k-1+\alpha}), \quad D^2 R \sim O(|\theta|^{k-2+\alpha})$$

with $\alpha \in (0, 1)$. It is a simple fact that $\{(\theta_1, \theta_2) | D^2 P = 0\} = \{(0, 0)\}$ and consequently $\{(\theta_1, \theta_2) | D^2 H = 0\} = \{(0, 0)\}$. Hence p is an isolated zero point of $D^2 u$. Therefore $S(u)$ consists of finitely many points in this case. $\square$

We consider the surface Σ parametrized by $\nabla u : \mathbb{S}^2 \rightarrow \mathbb{R}^3$. From (3.1), it follows that the Hessian always has one zero eigenvalue. Let $\lambda_1(x)$ and $\lambda_2(x)$ be the other two eigenvalues of $D^2 u(x)$. Because of equation (1.1), $\lambda_1(x) \lambda_2(x) < 0$ for all $x \in \mathbb{S}^2 \setminus S(u)$.

Lemma 3.2. *For any $x \in \mathbb{S}^2 \setminus S(u)$, the surface Σ is $C^{2,\alpha}$ at $\nabla u(x)$ with a normal vector given by x and the two principle curvatures given by $-1/\lambda_1(x)$ and $-1/\lambda_2(x)$.*

Proof. We may assume $x = p = (0, 0, 1) \in \mathbb{S}^2 \setminus S(u)$. Then locally at $\nabla u(p)$, Σ can be represented by

$$(3.2) \quad \mathbf{F}(x_1, x_2) = \nabla u(x_1, x_2, \sqrt{1 - x_1^2 - x_2^2}).$$

By differentiating the identity $x \cdot \nabla u(x) = u(x)$ twice with respect to x_i, x_j for $i, j = 1, 2$, we obtain

$$(3.3) \quad u_{3i}(p) = 0, \quad u_{3ij}(p) = -u_{ij}(p).$$

By differentiating (3.2) and making use of (3.3), we have

$$\mathbf{F}_i(0) = (u_{i1}(p), u_{i2}(p), 0)$$

and

$$\mathbf{F}_{ij}^{(3)}(0) = -u_{ij}(p)$$

where $\mathbf{F}_{ij}^{(3)}$ denotes the third component of the vector $\mathbf{F}_{ij}$. Then we get

$$\mathbf{F}_1(0) \times \mathbf{F}_2(0) = (0, 0, (u_{11}u_{22} - u_{12}^2)(p)).$$

It is easy to see that $(u_{11}u_{22} - u_{12}^2)(p) = \lambda_1\lambda_2(p) \neq 0$. Hence $(0, 0, 1)$ is a normal vector for Σ at $\nabla u(p)$. We see that $\nabla u = G^{-1}$ near $(0, 0, 1)$, where G^{-1} is the inverse of the Gauss map of Σ . Thus Σ is $C^{2,\alpha}$ nearby. We also get the first and second fundamental forms of Σ at $\nabla u(p)$ as follows

$$I = (u_{11}^2 + u_{12}^2)dx_1^2 + 2u_{12}(u_{11} + u_{22})dx_1dx_2 + (u_{21}^2 + u_{22}^2)dx_2^2,$$

$$II = -u_{11}dx_1^2 - 2u_{12}dx_1dx_2 - u_{22}dx_2^2,$$

where all u_{ij} are evaluated at p . Therefore the two principle curvatures are $-1/\lambda_1$ and $-1/\lambda_2$. $\square$

Remark 3.3. *In computing the principle curvatures of Σ in terms of the eigenvalues of D^2u , we differentiate u three times. By approximation, the conclusion still holds for $u \in C^{2,\alpha}$.*

Now we are ready for another proof of Theorem 1.1 in the $C^{2,\alpha}$ case.

Proof. We prove that $S(u) = \mathbb{S}^2$ by excluding the case (1) and (2) in Lemma 3.1. We assume $S(u)$ consists of at most finitely many points. First, the gradient surface Σ has supporting planes with normals along all the directions in $\mathbb{R}^3$. However, the saddle points in $\{\nabla u(x) \mid D^2u(x) \neq 0\}$ cannot support any supporting planes. Hence there are at most finitely many points on Σ with the supporting planes.

Claim. (Alexandrov) For any $x_0 \in S(u)$, the supporting plane to Σ at $\nabla u(x_0)$ can only have normal direction x_0 .

Therefore there are at most finitely many supporting planes to Σ . This is a contradiction. Now we prove the claim. Suppose there is another supporting plane P_1 to Σ at $\nabla u(x_0)$ with normal direction $x_1 \neq x_0$. Take a small neighborhood U of x_0 on $\mathbb{S}^2$ such that $D^2u(x) \neq 0$ for any $x \neq x_0 \in U$ and $U \cap S^* = \emptyset$, with S^* being a great circle through x_1 and $-x_1$. This can be done since x_0 is an isolated point in $S(u)$. Now lift P_1 along the x_1 direction to P so that $P \cap \nabla u(U) = C$ is a smooth close curve on P . We can take a point on C , say $\nabla u(x^*)$ with $x^* \in U$ and $x^* \neq x_0$, such that the normal to the plane curve C at $\nabla u(x^*)$ is along the intersection P and another plane through S^* . Then we see that the normal of the surface Σ at the regular

point $\nabla u(x^*)$ must be on S^* . On the other hand, the normal at $\nabla u(x^*)$ is $x^* \in U$. This contradiction completes the proof of the Claim.

By excluding the case (1) and (2) in Lemma 3.1, we are left with the case (3) $\mathcal{S}(u) = \mathbb{S}^2$. That is $D^2u \equiv 0$, and hence u is linear. This finishes the proof of Theorem 1.1 in the $C^{2,\alpha}$ case. $\square$

REFERENCES

- [1] Alexandrov, A. D., *Sur les théorèmes d'unicité pour les surfaces fermées*, C. R. Acad. Sci. URSS, N.s. **22** (1939), 99–102.
- [2] Bers, L., *Local behavior of solution of general linear elliptic equations*, Comm. Pure Appl. Math., **8** (1955), 473–496.
- [3] Bers, L. and Nirenberg, L., *On a representation theorem for linear elliptic systems with discontinuous coefficients and its application*, Convegno Internazionale sulle Equazioni Derivate e Parziali, August 1954, 111–140.
- [4] Caffarelli, L. A., Crandall, M. G., Kocan, M., and Świech, A., *On viscosity solutions of fully nonlinear equations with measurable ingredients*, Comm. Pure Appl. Math., **49** (1996), 365–397.
- [5] Gilbarg, D. and Trudinger, N. S., *Elliptic Partial Differential Equations of Second Order*, 2ed edition, Springer-Verlag, New York, 1983.
- [6] Han, Q., *Schauder estimates for elliptic operators with applications to nodal sets*, J. Geom. Analysis, **10** (2000), 455–480.
- [7] Lawson, H. B. Jr. and Osserman, R., *Non-existence, non-uniqueness and irregularity of solutions to the minimal surfaces system*, Acta Math., **139** (1977), 1–17.
- [8] Nadirashvili, N., *Nonclassical solutions to fully nonlinear elliptic equations*, to appear.
- [9] Pogorelov, A. V., *Solution of a problem of A. D. Aleksandrov*, (Russian), Dokl. Akad. Nauk, **360** (1998), 317–319.
- [10] Safonov, M. V., *Unimprovability of estimates of Holder constants for solutions of linear elliptic equations with measurable coefficients*, Mat. Sb., **132** (1987), 272–288; English translation in Math. USSR Sb., **60** (1988), 269–281.
- [11] Safonov, M. V., *Nonlinear Elliptic Equations of the Second Order*, Lecture notes, Univ. di Firenze, 1991.

DEPARTMENT OF MATHEMATICS, NOTRE DAME UNIVERSITY, NOTRE DAME, IN 46556
E-mail address: qhan@nd.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, 5734 S. UNIVERSITY AVE., CHICAGO, IL 60637
E-mail address: nicholas@math.uchicago.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, 5734 S. UNIVERSITY AVE., CHICAGO, IL 60637, AND UNIVERSITY OF WASHINGTON, SEATTLE, WA 98195
E-mail address: yuan@math.uchicago.edu